

WHAT IS CLAIMED IS:

1. A method for the bioproduction of C₆ to C₂₂ mono- and di-carboxylic acids comprising
- contacting, under aerobic conditions, a transformed *Pichia pastoris* characterized by a genetically-engineered alkane hydroxylating activity with at least one C₆ to C₂₂ straight chain hydrocarbon; and
 - recovering the C₆ to C₂₂ mono- and di-carboxylic acids.
2. The method of Claim 1 wherein the transformed *Pichia pastoris* is strain SW 64/65 identified as ATCC 74409; the at least one C₆ to C₂₂ straight chain hydrocarbon is dodecane; and the product recovered is dodecanedioic acid.
3. A transformed *Pichia pastoris* comprising
- at least one copy of a foreign gene encoding cytochrome P450 monooxygenase; and, optionally,
 - at least one copy of a foreign gene encoding cytochrome P450 reductase,
- each gene operably linked to suitable regulatory elements such that alkane hydroxylating activity is enhanced upon contact with at least one C₆ to C₂₂ straight chain hydrocarbon.
4. The transformed *Pichia pastoris* of Claim 3 wherein the foreign gene encoding cytochrome P450 monooxygenase is selected from the group consisting of Alk1-A (D12475), Alk2-A (X55881), Alk3-A (X55881), Alk4-A (D12716), Alk5-A (D12717), Alk6-A (D12718), Alk7 (D12719), and Alk8 (D12719).
5. The transformed *Pichia pastoris* of Claim 3 wherein the foreign gene encoding cytochrome P450 reductase is cytochrome P450 reductase (D25327).
6. A transformed *Pichia pastoris* strain characterized by an enhanced alkane hydroxylating activity and comprising,
- at least one DNA fragment from *Candida maltosa* ATCC 90677 selected from the group of DNA fragments encoding cytochrome P450 monooxygenase Alk1-A and cytochrome P450 monooxygenase Alk3-A; and, optionally,
 - at least one DNA fragment from *Candida maltosa* ATCC 90677 encoding cytochrome P450 reductase,
- each DNA fragment operably linked to suitable regulatory elements such that alkane hydroxylating activity is enhanced upon contact with at least one C₆ to C₂₂ straight chain hydrocarbon.
7. A transformed *Pichia pastoris* strain SW64/65 identified as ATCC 74409.
8. A method for the enhanced bioproduction of C₆ to C₂₂ mono- and di-carboxylic acids comprising

a) contacting, under aerobic conditions, a transformed *Candida maltosa* characterized by a genetically-engineered, enhanced alkane hydroxylating activity with at least one C₆ to C₂₂ straight chain hydrocarbon; and

b) recovering the C₆ to C₂₂ mono- and di-carboxylic acids.

5 9. The method of Claim 8 wherein

a) the genetically-engineered, enhanced alkane hydroxylating activity arises from

i) at least one additional copy of the genes encoding cytochrome P450 monooxygenase selected from the group consisting of Alk1-A (D12475), Alk2-A (X55881), Alk3-A (X55881), Alk4-A (D12716), Alk5-A (D12717), Alk6-A (D12718), Alk7 (D12719), and Alk8 (D12719); or

ii) at least one additional copy of the gene encoding cytochrome P450 reductase (D25327); or

iii) at least one additional copy of both the genes of i) and ii);

15 b) the at least one C₆ to C₂₂ straight chain hydrocarbon is dodecane;

and

c) the product recovered is dodecanedioic acid.

10. A transformed *Candida maltosa* comprising

a) at least one additional copy of a gene encoding cytochrome P450 monooxygenase; or

20 b) at least one additional copy of a gene encoding cytochrome P450 reductase; or

c) at least one additional copy of both the gene encoding P450 monooxygenase and the gene encoding cytochrome P450 reductase,

25 each gene operably linked to suitable regulatory elements such that alkane hydroxylating activity is enhanced upon contact with at least one C₆ to C₂₂ straight chain hydrocarbon.

11. The transformed *Candida maltosa* of Claim 10 wherein the genes encoding cytochrome P450 monooxygenase are selected from the group consisting of Alk1-A (D12475), Alk2-A (X55881), Alk3-A (X55881), Alk4-A (D12716), Alk5-A (D12717), Alk6-A (D12718), Alk7 (D12719), and Alk8 (D12719).

12. The transformed *Candida maltosa* of Claim 10 wherein the gene encoding cytochrome P450 reductase is cytochrome P450 reductase (D25327).

13. A transformed *Candida maltosa* strain comprising

a) at least one DNA fragment from *Candida maltosa* (ATCC 90677) selected from the group of DNA fragments encoding cytochrome P450 monooxygenase ALK1-A and cytochrome P450 monooxygenase ALK3-A, and

b) at least one DNA fragment from *Candida maltosa* (ATCC 90677) encoding cytochrome P450 reductase, each gene operably linked to suitable regulatory elements such that alkane hydroxylating activity is enhanced upon contact with at least one C₆ to C₂₂ straight chain hydrocarbon.

14. A method for the enhanced bioproduction of C₆ to C₂₂ mono- and di-carboxylic acids comprising

a) contacting, under aerobic conditions, transformed *Candida maltosa* characterized by a genetically-engineered, blocked β -oxidation pathway with at least one C₆ to C₂₂ straight chain hydrocarbon; and

b) recovering the C₆ to C₂₂ mono- and di-carboxylic acids.

15. The method of Claim 14 wherein the transformed *Candida maltosa* β -oxidation pathway is functionally blocked by disruption of both POX4 genes encoding acyl-CoA oxidase.

16. A transformed *Candida maltosa* characterized by a β -oxidation pathway functionally blocked by disruption of both POX4 genes encoding acyl-CoA oxidase.

17. A transformed *Candida maltosa* characterized by a β -oxidation pathway functionally blocked by disruption of both POX4 genes encoding acyl-CoA oxidase using a single URA3 selectable marker.

18. A transformed *Candida maltosa* strain SW81/82 identified as ATCC 74431.

19. A method for the enhanced bioproduction of C₆ to C₂₂ mono- and di-carboxylic acids comprising

a) contacting, under aerobic conditions, transformed *Candida maltosa* characterized by

i) a genetically-engineered, enhanced alkane hydroxylating activity, and

ii) a genetically-engineered, blocked β -oxidation pathway, with at least one C₆ to C₂₂ straight chain hydrocarbon; and

b) recovering the C₆ to C₂₂ mono- and di-carboxylic acids.

20. A transformed *Candida maltosa* characterized by

a) an enhanced alkane hydroxylating activity arising from

i) at least one additional copy of a gene encoding cytochrome P450 monooxygenase selected from the group consisting of Alk1-A (D12475), Alk2-A (X55881), Alk3-A (X55881), Alk4-A (D12716), Alk5-A (D12717), Alk6-A (D12718), Alk7 (D12719), and Alk8 (D12719), or

ii) at least one additional copy of a gene encoding cytochrome P450 reductase (D25327), or

Sub
A5

iii) at least one additional copy of both the genes i) and ii); and
b) a β -oxidation pathway functionally blocked by disruption of both
POX4 genes encoding acyl-CoA oxidase.

Sub
A6

21. The transformed *Candida maltosa* strain of Claim 20 wherein the
enhanced alkane hydroxylating activity of a) arises from DNA fragments
encoding cytochrome P450 monooxygenase ALK1-A and cytochrome P450
monooxygenase ALK3-A.

22. A transformed *Candida maltosa* strain SW84/87.2 identified as
ATCC 74430.

23. The method of Claims 1, 8, 14, or 19 wherein the at least one C₆ to
C₂₂ straight chain hydrocarbon is selected from the group consisting of hexane,
heptane, octane, nonane, decane, undecane, dodecane, tridecane, tetradecane,
pentadecane, hexadecane, heptadecane, octadecane, nonadecane, eicosane,
reneicosane, docosane and their respective mono-carboxylic acids and esters.

24. A DNA fragment comprising a) a first *Candida maltosa* promoter
operably linked to DNA encoding at least one polypeptide from *Candida maltosa*
and b) a second *Candida maltosa* promoter operably linked to DNA encoding at
least one polypeptide from *Candida maltosa*.

25. A DNA fragment comprising a) a first *Candida maltosa* promoter
operably linked to a gene encoding a *Candida maltosa* cytochrome P450
monooxygenase and b) a second *Candida maltosa* promoter operably linked to a
gene encoding a *Candida maltosa* cytochrome P450 reductase.

26. A DNA fragment comprising a) a first *Candida maltosa* PGK
promoter which is operably linked to a gene encoding cytochrome P450
monooxygenase selected from the group consisting of Alk1-A (D12475), Alk2-A
(X55881), Alk3-A (X55881), Alk4-A (D12716), Alk5-A (D12717), Alk6-A
(D12718), Alk7 (D12719), and Alk8 (D12719) and b) a second *Candida maltosa*
PGK promoter operably linked to a gene encoding a *Candida maltosa* cytochrome
P450 reductase.

27. A plasmid selected from the group consisting of pSW84 and pSW87.

Sub
A7

09116502.071698